

Hindawi Publishing Corporation
Fixed Point Theory and Applications
Volume 2007, Article ID 29091, 10 pages
doi:10.1155/2007/29091

Research Article

Iteration Scheme with Perturbed Mapping for Common Fixed Points of a Finite Family of Nonexpansive Mappings

Yeong-Cheng Liou, Yonghong Yao, and Rudong Chen

Received 17 December 2006; Revised 6 February 2007; Accepted 6 February 2007

Recommended by H  l  ne Frankowska

We propose an iteration scheme with perturbed mapping for approximation of common fixed points of a finite family of nonexpansive mappings $\{T_i\}_{i=1}^N$. We show that the proposed iteration scheme converges to the common fixed point $x^* \in \bigcap_{i=1}^N \text{Fix}(T_i)$ which solves some variational inequality.

Copyright   2007 Yeong-Cheng Liou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction and preliminaries

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, respectively. A mapping T with domain $D(T)$ and range $R(T)$ in H is called nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in D(T). \quad (1.1)$$

Let $\{T_i\}_{i=1}^N$ be a finite family of nonexpansive self-maps of H . Denote the common fixed points set of $\{T_i\}_{i=1}^N$ by $\bigcap_{i=1}^N \text{Fix}(T_i)$. Let $F : H \rightarrow H$ be a mapping such that for some constants $k, \eta > 0$, F is k -Lipschitzian and η -strongly monotone. Let $\{\alpha_n\}_{n=1}^\infty \subset (0, 1)$, $\{\lambda_n\}_{n=1}^\infty \subset [0, 1)$ and take a fixed number $\mu \in (0, 2\eta/k^2)$. The iterative schemes concerning nonlinear operators have been studied extensively by many authors, you may refer to [1–12]. Especially, in [13], Zeng and Yao introduced the following implicit iteration process with perturbed mapping F .

For an arbitrary initial point $x_0 \in H$, the sequence $\{x_n\}_{n=1}^\infty$ is generated as follows:

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) [T_n x_n - \lambda_n \mu F(T_n x_n)], \quad n \geq 1, \quad (1.2)$$

where $T_n := T_{n \bmod N}$.

2 Fixed Point Theory and Applications

Using this iteration process, they proved the following weak and strong convergence theorems for nonexpansive mappings in Hilbert spaces.

THEOREM 1.1 (see [13]). *Let H be a real Hilbert space and let $F : H \rightarrow H$ be a mapping such that for some constants $k, \eta > 0$, F is k -Lipschitzian and η -strongly monotone. Let $\{T_i\}_{i=1}^N$ be N nonexpansive self-mappings of H such that $\bigcap_{i=1}^N \text{Fix}(T_i) \neq \emptyset$. Let $\mu \in (0, 2\eta/k^2)$ and $x_0 \in H$. Let $\{\lambda_n\}_{n=1}^\infty \subset [0, 1)$ and $\{\alpha_n\}_{n=1}^\infty \subset (0, 1)$ satisfying the conditions $\sum_{n=1}^\infty \lambda_n < \infty$ and $\alpha \leq \alpha_n \leq \beta$, $n \geq 1$, for some $\alpha, \beta \in (0, 1)$. Then the sequence $\{x_n\}_{n=1}^\infty$ defined by (1.2) converges weakly to a common fixed point of the mappings $\{T_i\}_{i=1}^N$.*

THEOREM 1.2 (see [13]). *Let H be a real Hilbert space and let $F : H \rightarrow H$ be a mapping such that for some constants $k, \eta > 0$, F is k -Lipschitzian and η -strongly monotone. Let $\{T_i\}_{i=1}^N$ be N nonexpansive self-mappings of H such that $\bigcap_{i=1}^N \text{Fix}(T_i) \neq \emptyset$. Let $\mu \in (0, 2\eta/k^2)$ and $x_0 \in H$. Let $\{\lambda_n\}_{n=1}^\infty \subset [0, 1)$ and $\{\alpha_n\}_{n=1}^\infty \subset (0, 1)$ satisfying the conditions $\sum_{n=1}^\infty \lambda_n < \infty$ and $\alpha \leq \alpha_n \leq \beta$, $n \geq 1$, for some $\alpha, \beta \in (0, 1)$. Then the sequence $\{x_n\}_{n=1}^\infty$ defined by (1.2) converges strongly to a common fixed point of the mappings $\{T_i\}_{i=1}^N$ if and only if*

$$\liminf_{n \rightarrow \infty} d\left(x_n, \bigcap_{i=1}^N \text{Fix}(T_i)\right) = 0. \quad (1.3)$$

Very recently, Wang [14] considered an explicit iterative scheme with perturbed mapping F and obtained the following result.

THEOREM 1.3. *Let H be a Hilbert space, let $T : H \rightarrow H$ be a nonexpansive mapping with $F(T) \neq \emptyset$, and let $F : H \rightarrow H$ be an η -strongly monotone and k -Lipschitzian mapping. For any given $x_0 \in H$, $\{x_n\}$ is defined by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 0, \quad (1.4)$$

where $T^{\lambda_{n+1}} x_n = T x_n - \lambda_{n+1} \mu F(T x_n)$, $\{\alpha_n\}$ and $\{\lambda_n\} \subset [0, 1)$ satisfy the following conditions:

- (1) $\alpha \leq \alpha_n \leq \beta$ for some $\alpha, \beta \in (0, 1)$;
- (2) $\sum_{n=1}^\infty \lambda_n < \infty$;
- (3) $0 < \mu < 2\eta/k^2$.

Then

- (1) $\{x_n\}$ converges weakly to a fixed point of T ,
- (2) $\{x_n\}$ converges strongly to a fixed point of T if and only if

$$\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0. \quad (1.5)$$

This naturally brings us the following questions.

Questions 1.4. Let $T_i : H \rightarrow H$ ($i = 1, 2, \dots, N$) be a finite family of nonexpansive mappings and F is k -Lipschitzian and η -strongly monotone.

- (i) Could we construct an explicit iterative algorithm to approximate the common fixed points of the mappings $\{T_i\}_{i=1}^N$?
- (ii) Could we remove the assumption (2) imposed on the sequence $\{x_n\}$?

Motivated and inspired by the above research work of Zeng and Yao [13] and Wang [14], in this paper, we will propose a new explicit iteration scheme with perturbed mapping for approximation of common fixed points of a finite family of nonexpansive self-mappings of H . We will establish strong convergence theorem for this explicit iteration scheme. To be more specific, let $\alpha_{n1}, \alpha_{n2}, \dots, \alpha_{nN} \in (0, 1]$, $n \in N$. Given the mappings T_1, T_2, \dots, T_N , following [15], one can define, for each n , mappings $U_{n1}, U_{n2}, \dots, U_{nN}$ by

$$\begin{aligned} U_{n1} &= \alpha_{n1} T_1 + (1 - \alpha_{n1}) I, \\ U_{n2} &= \alpha_{n2} T_2 U_{n1} + (1 - \alpha_{n2}) I, \\ &\vdots \\ U_{n,N-1} &= \alpha_{n,N-1} T_{N-1} U_{n,N-2} + (1 - \alpha_{n,N-1}) I, \\ W_n &:= U_{nN} = \alpha_{nN} T_N U_{n,N-1} + (1 - \alpha_{nN}) I. \end{aligned} \quad (1.6)$$

Such a mapping W_n is called the W -mapping generated by T_1, \dots, T_N and $\alpha_{n1}, \dots, \alpha_{nN}$.

First we introduce the following explicit iteration scheme with perturbed mapping F .

For an arbitrary initial point $x_0 \in H$, the sequence $\{x_n\}_{n=1}^\infty$ is generated iteratively by

$$x_{n+1} = \beta x_n + (1 - \beta) [W_n x_n - \lambda_n \mu F(W_n x_n)], \quad n \geq 0, \quad (1.7)$$

where $\{\lambda_n\}$ is a sequence in $(0, 1)$, β is a constant in $(0, 1)$, F is k -Lipschitzian and η -strongly monotone, and W_n is the W -mapping defined by (1.6).

We have the following crucial conclusion concerning W_n .

PROPOSITION 1.5 (see [15]). *Let C be a nonempty closed convex subset of a Banach space E . Let T_1, T_2, \dots, T_N be nonexpansive mappings of C into itself such that $\bigcap_{i=1}^N \text{Fix}(T_i)$ is nonempty, and let $\alpha_{n1}, \alpha_{n2}, \dots, \alpha_{nN}$ be real numbers such that $0 < \alpha_{ni} \leq b < 1$ for any $i \in N$. For any $n \in N$, let W_n be the W -mapping of C into itself generated by T_N, T_{N-1}, \dots, T_1 and $\alpha_{nN}, \alpha_{n,N-1}, \dots, \alpha_{n1}$. Then W_n is nonexpansive. Further, if E is strictly convex, then $\text{Fix}(W_n) = \bigcap_{i=1}^N \text{Fix}(T_i)$.*

Now we recall some basic notations. Let $T : H \rightarrow H$ be nonexpansive mapping and $F : H \rightarrow H$ be a mapping such that for some constants $k, \eta > 0$, F is k -Lipschitzian and η -strongly monotone; that is, F satisfies the following conditions:

$$\begin{aligned} \|Fx - Fy\| &\leq k\|x - y\|, \quad \forall x, y \in H, \\ \langle Fx - Fy, x - y \rangle &\geq \eta\|x - y\|^2, \quad \forall x, y \in H, \end{aligned} \quad (1.8)$$

respectively. We may assume, without loss of generality, that $\eta \in (0, 1)$ and $k \in [1, \infty)$. Under these conditions, it is well known that the variational inequality problem—find $x^* \in \bigcap_{i=1}^N \text{Fix}(T_i)$ such that

$$VI \left(F, \bigcap_{i=1}^N \text{Fix}(T_i) \right) : \langle F(x^*), x - x^* \rangle \geq 0, \quad \forall x \in \bigcap_{i=1}^N \text{Fix}(T_i), \quad (1.9)$$

4 Fixed Point Theory and Applications

has a unique solution $x^* \in \bigcap_{i=1}^N \text{Fix}(T_i)$. [Note: the unique existence of the solution $x^* \in \bigcap_{i=1}^N \text{Fix}(T_i)$ is guaranteed automatically because F is k -Lipschitzian and η -strongly monotone over $\bigcap_{i=1}^N \text{Fix}(T_i)$.]

For any given numbers $\lambda \in [0, 1]$ and $\mu \in (0, 2\eta/k^2)$, we define the mapping $T^\lambda : H \rightarrow H$ by

$$T^\lambda x := Tx - \lambda \mu F(Tx), \quad \forall x \in H. \quad (1.10)$$

Concerning the corresponding result of $T^\lambda x$, you can find it in [16].

LEMMA 1.6 (see [16]). *If $0 \leq \lambda < 1$ and $0 < \mu < 2\eta/k^2$, then there holds for $T^\lambda : H \rightarrow H$,*

$$\|T^\lambda x - T^\lambda y\| \leq (1 - \lambda\tau)\|x - y\|, \quad \forall x, y \in H, \quad (1.11)$$

where $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu k^2)} \in (0, 1)$.

Next, let us state four preliminary results which will be needed in the sequel. Lemma 1.7 is very interesting and important, you may find it in [17], the original prove can be found in [18]. Lemmas 1.8 and 1.9 well-known demiclosedness principle and subdifferential inequality, respectively. Lemma 1.10 is basic and important result, please consult it in [19].

LEMMA 1.7 (see [17]). *Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space X and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with*

$$0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1. \quad (1.12)$$

Suppose

$$x_{n+1} = (1 - \beta_n)y_n + \beta_n x_n, \quad (1.13)$$

for all integers $n \geq 0$ and

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (1.14)$$

Then, $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.

LEMMA 1.8 (see [20]). *Assume that T is a nonexpansive self-mapping of a closed convex subset C of a Hilbert space H . If T has a fixed point, then $I - T$ is demiclosed. That is, whenever $\{x_n\}$ is a sequence in C weakly converging to some $x \in C$ and the sequence $\{(I - T)x_n\}$ strongly converges to some y , it follows that $(I - T)x = y$. Here, I is the identity operator of H .*

LEMMA 1.9 (see [21]). $\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle$ for all $x, y \in H$.

LEMMA 1.10 (see [19]). *Assume that $\{a_n\}$ is a sequence of nonnegative real numbers such that*

$$a_{n+1} \leq (1 - \gamma_n)a_n + \delta_n, \quad (1.15)$$

where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence such that

- (1) $\sum_{n=1}^{\infty} \gamma_n = \infty$,
- (2) $\limsup_{n \rightarrow \infty} \delta_n / \gamma_n \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n| < \infty$.

Then $\lim_{n \rightarrow \infty} a_n = 0$.

2. Main result

Now we state and prove our main result.

THEOREM 2.1. *Let H be a real Hilbert space and let $F : H \rightarrow H$ be a k -Lipschitzian and η -strongly monotone mapping. Let $\{T_i\}_{i=1}^N$ be a finite family of nonexpansive self-mappings of H such that $\bigcap_{i=1}^N \text{Fix}(T_i) \neq \emptyset$. Let $\mu \in (0, 2\eta/k^2)$. Suppose the sequences $\{\alpha_{n,i}\}_{i=1}^N$ satisfy $\lim_{n \rightarrow \infty} (\alpha_{n,i} - \alpha_{n-1,i}) = 0$, for all $i = 1, 2, \dots, N$. If $\{\lambda_n\}_{n=1}^{\infty} \subset [0, 1)$ satisfy the following conditions:*

- (i) $\lim_{n \rightarrow \infty} \lambda_n = 0$;
- (ii) $\sum_{n=0}^{\infty} \lambda_n = \infty$,

then the sequence $\{x_n\}_{n=1}^{\infty}$ defined by (1.7) converges strongly to a common fixed point $x^ \in \bigcap_{i=1}^N \text{Fix}(T_i)$ which solves the variational inequality (1.9).*

Proof. Let x^* be an arbitrary element of $\bigcap_{i=1}^N \text{Fix}(T_i)$. Observe that

$$\begin{aligned} \|x_{n+1} - x^*\| &= \|\beta x_n + (1 - \beta)W_n^{\lambda_n} x_n - x^*\| \\ &\leq \beta \|x_n - x^*\| + (1 - \beta) \|W_n^{\lambda_n} x_n - x^*\|, \end{aligned} \quad (2.1)$$

where $W_n^{\lambda_n} x := W_n x - \lambda_n \mu F(W_n x)$. Note that

$$W_n^{\lambda_n} x^* = x^* - \lambda_n \mu F(x^*). \quad (2.2)$$

Utilizing Lemma 1.6, we have

$$\begin{aligned} \|W_n^{\lambda_n} x_n - x^*\| &= \|W_n^{\lambda_n} x_n - W_n^{\lambda_n} x^* + W_n^{\lambda_n} x^* - x^*\| \\ &\leq \|W_n^{\lambda_n} x_n - W_n^{\lambda_n} x^*\| + \|W_n^{\lambda_n} x^* - x^*\| \\ &\leq (1 - \lambda_n \tau) \|x_n - x^*\| + \lambda_n \mu \|F(x^*)\|. \end{aligned} \quad (2.3)$$

From (2.1) and (2.3), we have

$$\begin{aligned} \|x_{n+1} - x^*\| &\leq [\beta + (1 - \beta)(1 - \lambda_n \tau)] \|x_n - x^*\| + (1 - \beta) \lambda_n \mu \|F(x^*)\| \\ &= [1 - (1 - \beta) \lambda_n \tau] \|x_n - x^*\| + (1 - \beta) \lambda_n \mu \|F(x^*)\| \\ &\leq \max \left\{ \|x_0 - x^*\|, \left(\frac{\mu}{\tau} \right) \|F(x^*)\| \right\}. \end{aligned} \quad (2.4)$$

Hence, $\{x_n\}$ is bounded. We also can obtain that $\{W_n x_n\}$, $\{T_i U_{n,j} x_n\}$ ($i = 1, \dots, N$; $j = 1, \dots, N$), and $\{F(W_n x_n)\}$ are all bounded.

We will use M to denote the possible different constants appearing in the following reasoning.

6 Fixed Point Theory and Applications

We note that

$$\begin{aligned}
 & \|W_{n+1}^{\lambda_{n+1}}x_{n+1} - W_n^{\lambda_n}x_n\| \\
 &= \|W_{n+1}x_{n+1} - W_nx_n - \lambda_{n+1}\mu F(W_{n+1}x_{n+1}) + \lambda_n\mu F(W_nx_n)\| \\
 &\leq \|W_{n+1}x_{n+1} - W_nx_n\| + \lambda_{n+1}\mu \|F(W_{n+1}x_{n+1})\| + \lambda_n\mu \|F(W_nx_n)\| \\
 &\leq \|W_{n+1}x_{n+1} - W_{n+1}x_n\| + \|W_{n+1}x_n - W_nx_n\| + (\lambda_{n+1} + \lambda_n)M \\
 &\leq \|x_{n+1} - x_n\| + \|W_{n+1}x_n - W_nx_n\| + (\lambda_{n+1} + \lambda_n)M.
 \end{aligned} \tag{2.5}$$

From (1.6), since T_N and $U_{n,N}$ are nonexpansive,

$$\begin{aligned}
 & \|W_{n+1}x_n - W_nx_n\| \\
 &= \|\alpha_{n+1,N}T_NU_{n+1,N-1}x_n + (1 - \alpha_{n+1,N})x_n - \alpha_{n,N}T_NU_{n,N-1}x_n - (1 - \alpha_{n,N})x_n\| \\
 &\leq \|\alpha_{n+1,N}T_NU_{n+1,N-1}x_n - \alpha_{n,N}T_NU_{n,N-1}x_n\| + |\alpha_{n+1,N} - \alpha_{n,N}| \|x_n\| \\
 &\leq \|\alpha_{n+1,N}(T_NU_{n+1,N-1}x_n - T_NU_{n,N-1}x_n)\| + |\alpha_{n+1,N} - \alpha_{n,N}| \|T_NU_{n,N-1}x_n\| \\
 &\quad + |\alpha_{n+1,N} - \alpha_{n,N}| \|x_n\| \\
 &\leq \alpha_{n+1,N} \|U_{n+1,N-1}x_n - U_{n,N-1}x_n\| + 2M |\alpha_{n+1,N} - \alpha_{n,N}|.
 \end{aligned} \tag{2.6}$$

Again, from (1.6), we have

$$\begin{aligned}
 & \|U_{n+1,N-1}x_n - U_{n,N-1}x_n\| \\
 &= \|\alpha_{n+1,N-1}T_{N-1}U_{n+1,N-2}x_n + (1 - \alpha_{n+1,N-1})x_n \\
 &\quad - \alpha_{n,N-1}T_{N-1}U_{n,N-2}x_n - (1 - \alpha_{n,N-1})x_n\| \\
 &\leq \|\alpha_{n+1,N-1}T_{N-1}U_{n+1,N-2}x_n - \alpha_{n,N-1}T_{N-1}U_{n,N-2}x_n\| \\
 &\quad + |\alpha_{n+1,N-1} - \alpha_{n,N-1}| \|x_n\| \\
 &\leq |\alpha_{n+1,N-1} - \alpha_{n,N-1}| \|x_n\| + |\alpha_{n+1,N-1} - \alpha_{n,N-1}| M \\
 &\quad + \alpha_{n+1,N-1} \|T_{N-1}U_{n+1,N-2}x_n - T_{N-1}U_{n,N-2}x_n\| \\
 &\leq 2M |\alpha_{n+1,N-1} - \alpha_{n,N-1}| + \alpha_{n+1,N-1} \|U_{n+1,N-2}x_n - U_{n,N-2}x_n\| \\
 &\leq 2M |\alpha_{n+1,N-1} - \alpha_{n,N-1}| + \|U_{n+1,N-2}x_n - U_{n,N-2}x_n\|.
 \end{aligned} \tag{2.7}$$

Therefore, we have

$$\begin{aligned}
& \|U_{n+1,N-1}x_n - U_{n,N-1}x_n\| \\
& \leq 2M|\alpha_{n+1,N-1} - \alpha_{n,N-1}| + 2M|\alpha_{n+1,N-2} - \alpha_{n,N-2}| \\
& \quad + \|U_{n+1,N-3}x_n - U_{n,N-3}x_n\| \\
& \leq 2M \sum_{i=2}^{N-1} |\alpha_{n+1,i} - \alpha_{n,i}| + \|U_{n+1,1}x_n - U_{n,1}x_n\| \\
& = \|\alpha_{n+1,1}T_1x_n + (1 - \alpha_{n+1,1})x_n - \alpha_{n,1}T_1x_n - (1 - \alpha_{n,1})x_n\| \\
& \quad + 2M \sum_{i=2}^{N-1} |\alpha_{n+1,i} - \alpha_{n,i}|,
\end{aligned} \tag{2.8}$$

then

$$\begin{aligned}
& \|U_{n+1,N-1}x_n - U_{n,N-1}x_n\| \\
& \leq |\alpha_{n+1,1} - \alpha_{n,1}| \|x_n\| + \|\alpha_{n+1,1}T_1x_n - \alpha_{n,1}T_1x_n\| \\
& \quad + 2M \sum_{i=2}^{N-1} |\alpha_{n+1,i} - \alpha_{n,i}| \leq 2M \sum_{i=1}^{N-1} |\alpha_{n+1,i} - \alpha_{n,i}|.
\end{aligned} \tag{2.9}$$

Substituting (2.9) into (2.6), we have

$$\begin{aligned}
\|W_{n+1}x_n - W_nx_n\| & \leq 2M|\alpha_{n+1,N} - \alpha_{n,N}| + 2\alpha_{n+1,N}M \sum_{i=1}^{N-1} |\alpha_{n+1,i} - \alpha_{n,i}| \\
& \leq 2M \sum_{i=1}^N |\alpha_{n+1,i} - \alpha_{n,i}|.
\end{aligned} \tag{2.10}$$

Substituting (2.10) into (2.5), we have

$$\|W_{n+1}^{\lambda_{n+1}}x_{n+1} - W_n^{\lambda_n}x_n\| \leq \|x_{n+1} - x_n\| + 2M \sum_{i=1}^N |\alpha_{n+1,i} - \alpha_{n,i}| + (\lambda_{n+1} + \lambda_n)M, \tag{2.11}$$

which implies that

$$\limsup_{n \rightarrow \infty} (\|W_{n+1}^{\lambda_{n+1}}x_{n+1} - W_n^{\lambda_n}x_n\| - \|x_{n+1} - x_n\|) \leq 0. \tag{2.12}$$

We note that $x_{n+1} = \beta x_n + (1 - \beta)W_n^{\lambda_n}x_n$ and $0 < \beta < 1$, then from Lemma 1.7 and (2.12), we have $\lim_{n \rightarrow \infty} \|W_n^{\lambda_n}x_n - x_n\| = 0$. It follows that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} (1 - \beta) \|W_n^{\lambda_n}x_n - x_n\| = 0. \tag{2.13}$$

On the other hand,

$$\begin{aligned}
\|x_n - W_nx_n\| & \leq \|x_{n+1} - x_n\| + \|x_{n+1} - W_nx_n\| \\
& \leq \|x_{n+1} - x_n\| + \beta \|x_n - W_nx_n\| + (1 - \beta)\lambda_n\mu \|F(W_nx_n)\|,
\end{aligned} \tag{2.14}$$

8 Fixed Point Theory and Applications

that is,

$$\|x_n - W_n x_n\| \leq \frac{1}{1-\beta} \|x_{n+1} - x_n\| + \lambda_n \mu \|F(W_n x_n)\|, \quad (2.15)$$

this together with (i) and (2.13) imply

$$\lim_{n \rightarrow \infty} \|x_n - W_n x_n\| = 0. \quad (2.16)$$

We next show that

$$\limsup_{n \rightarrow \infty} \langle -F(x^*), x_n - x^* \rangle \leq 0. \quad (2.17)$$

To prove this, we pick a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle -F(x^*), x_n - x^* \rangle = \lim_{i \rightarrow \infty} \langle -F(x^*), x_{n_i} - x^* \rangle. \quad (2.18)$$

Without loss of generality, we may further assume that $x_{n_i} \rightarrow z$ weakly for some $z \in H$.

By Lemma 1.8 and (2.16), we have

$$z \in \text{Fix}(W_n), \quad (2.19)$$

this together with Proposition 1.5 imply that

$$z \in \bigcap_{i=1}^N \text{Fix}(T_i). \quad (2.20)$$

Since x^* solves the variational inequality (1.9), then we obtain

$$\limsup_{n \rightarrow \infty} \langle -F(x^*), x_n - x^* \rangle = \langle -F(x^*), z - x^* \rangle \leq 0. \quad (2.21)$$

Finally, we show that $x_n \rightarrow x^*$. Indeed, from Lemma 1.9, we have

$$\begin{aligned} & \|x_{n+1} - x^*\|^2 \\ &= \|\beta(x_n - x^*) + (1-\beta)(W_n^{\lambda_n} x_n - W_n^{\lambda_n} x^*) + (1-\beta)(W_n^{\lambda_n} x^* - x^*)\|^2 \\ &\leq \|\beta(x_n - x^*) + (1-\beta)(W_n^{\lambda_n} x_n - W_n^{\lambda_n} x^*)\|^2 + 2(1-\beta)\langle W_n^{\lambda_n} x^* - x^*, x_{n+1} - x^* \rangle \\ &\leq [\beta\|x_n - x^*\| + (1-\beta)\|W_n^{\lambda_n} x_n - W_n^{\lambda_n} x^*\|]^2 + 2(1-\beta)\lambda_n \mu \langle -F(x^*), x_{n+1} - x^* \rangle \\ &\leq [\beta\|x_n - x^*\| + (1-\beta)(1-\lambda_n \tau)\|x_n - x^*\|]^2 + 2(1-\beta)\lambda_n \mu \langle -F(x^*), x_{n+1} - x^* \rangle \\ &\leq [1 - (1-\beta)\tau\lambda_n]\|x_n - x^*\|^2 + (1-\beta)\tau\lambda_n \left\{ 2\frac{\mu}{\tau} \langle -F(x^*), x_{n+1} - x^* \rangle \right\}. \end{aligned} \quad (2.22)$$

Now applying Lemma 1.10 and (2.21) to (2.22) concludes that $x_n \rightarrow x^*$ ($n \rightarrow \infty$). This completes the proof. \square

Acknowledgments

The authors thank the referees for their suggestions and comments which led to the present version. The research was partially supposed by Grant NSC 95-2221-E-230-017.

References

- [1] L.-C. Zeng and J.-C. Yao, "Stability of iterative procedures with errors for approximating common fixed points of a couple of q -contractive-like mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 321, no. 2, pp. 661–674, 2006.
- [2] Y.-C. Lin, N.-C. Wong, and J.-C. Yao, "Strong convergence theorems of Ishikawa iteration process with errors for fixed points of Lipschitz continuous mappings in Banach spaces," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 543–552, 2006.
- [3] L.-C. Zeng, N.-C. Wong, and J.-C. Yao, "Strong convergence theorems for strictly pseudocontractive mappings of Browder-Petryshyn type," *Taiwanese Journal of Mathematics*, vol. 10, no. 4, pp. 837–849, 2006.
- [4] Y.-C. Lin, "Three-step iterative convergence theorems with errors in Banach spaces," *Taiwanese Journal of Mathematics*, vol. 10, no. 1, pp. 75–86, 2006.
- [5] L.-C. Zeng, G. M. Lee, and N.-C. Wong, "Ishikawa iteration with errors for approximating fixed points of strictly pseudocontractive mappings of Browder-Petryshyn type," *Taiwanese Journal of Mathematics*, vol. 10, no. 1, pp. 87–99, 2006.
- [6] S. Schaible, J.-C. Yao, and L.-C. Zeng, "A proximal method for pseudomonotone type variational-like inequalities," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 497–513, 2006.
- [7] L.-C. Zeng, L. J. Lin, and J.-C. Yao, "Auxiliary problem method for mixed variational-like inequalities," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 515–529, 2006.
- [8] L.-C. Zeng, N.-C. Wong, and J.-C. Yao, "On the convergence analysis of modified hybrid steepest-descent methods with variable parameters for variational inequalities," to appear in *Journal of Optimization Theory and Applications*.
- [9] L. C. Ceng, P. Cubiotti, and J.-C. Yao, "Approximation of common fixed points of families of nonexpansive mappings," to appear in *Taiwanese Journal of Mathematics*.
- [10] L. C. Ceng, P. Cubiotti, and J.-C. Yao, "Strong convergence theorems for finitely many nonexpansive mappings and applications," to appear in *Nonlinear Analysis*.
- [11] L.-C. Zeng, S. Y. Wu, and J.-C. Yao, "Generalized KKM theorem with applications to generalized minimax inequalities and generalized equilibrium problems," *Taiwanese Journal of Mathematics*, vol. 10, no. 6, pp. 1497–1514, 2006.
- [12] L. C. Ceng, C. Lee, and J.-C. Yao, "Strong weak convergence theorems of implicit hybrid steepest-descent methods for variational inequalities," to appear in *Taiwanese Journal of Mathematics*.
- [13] L.-C. Zeng and J.-C. Yao, "Implicit iteration scheme with perturbed mapping for common fixed points of a finite family of nonexpansive mappings," *Nonlinear Analysis*, vol. 64, no. 11, pp. 2507–2515, 2006.
- [14] L. Wang, "An iteration method for nonexpansive mappings in Hilbert spaces," *Fixed Point Theory and Applications*, vol. 2007, Article ID 28619, 8 pages, 2007.
- [15] W. Takahashi and K. Shimoji, "Convergence theorems for nonexpansive mappings and feasibility problems," *Mathematical and Computer Modelling*, vol. 32, no. 11–13, pp. 1463–1471, 2000.
- [16] H. K. Xu and T. H. Kim, "Convergence of hybrid steepest-descent methods for variational inequalities," *Journal of Optimization Theory and Applications*, vol. 119, no. 1, pp. 185–201, 2003.

- [17] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [18] T. Suzuki, "Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces," *Fixed Point Theory and Applications*, vol. 2005, no. 1, pp. 103–123, 2005.
- [19] H.-K. Xu, "Viscosity approximation methods for nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 298, no. 1, pp. 279–291, 2004.
- [20] K. Geobel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, UK, 1990.
- [21] G. Marino and H.-K. Xu, "Convergence of generalized proximal point algorithms," *Communications on Pure and Applied Analysis*, vol. 3, no. 4, pp. 791–808, 2004.

Yeong-Cheng Liou: Department of Information Management, Cheng Shiu University,
Kaohsiung 833, Taiwan
Email address: simplex_liou@hotmail.com

Yonghong Yao: Department of Mathematics, Tianjin Polytechnic University, Tianji 300160, China
Email address: yuyanrong@tjpu.edu.cn

Rudong Chen: Department of Mathematics, Tianjin Polytechnic University, Tianji 300160, China
Email address: chenrd@tjpu.edu.cn